

THE GEOMETRY OF SYSTEMS OF THIRD ORDER DIFFERENTIAL EQUATIONS INDUCED BY SECOND ORDER LAGRANGIANS

IOAN BUCATARU AND RADU MIRON

ABSTRACT. A dynamical system on the total space of the fibre bundle of second order accelerations, T^2M , is defined as a third order vector field S on T^2M , called semispray, which is mapped by the second order tangent structure into one of the Liouville vector field. For a regular Lagrangian of second order we prove that this semispray is uniquely determined by two associated Cartan-Poincaré one-forms. To study the geometry of this semispray we construct a nonlinear connection, which is a Lagrangian subbundle for the presymplectic structure. Using this semispray and the associated nonlinear connection we define covariant derivatives of first and second order. With respect to this, the second order dynamical derivative of the Lagrangian metric tensor vanishes.

2000 MSC: 53B40, 53C60, 70G45

Keywords and phrases: Craig-Synge equations, third order vector field, semispray, nonlinear connection, second order Lagrangian.

1. INTRODUCTION

The notion of spray as a vector field that lives on the total space of the first order tangent bundle was introduced by Ambrose et al. [2]. For a regular Lagrangian of order one, solutions of the Euler-Lagrange equations are integral curves of the canonical semispray, which is a second order vector field. The canonical semispray of a Lagrange space is uniquely determined by its symplectic structure and the energy of the space. This point of view allows for studying first integrals, Cartan symmetries, and Noether type theorems and it has been investigated by Klein [24], Godbillon [20], Abraham and Marsden [1], Arnold [4], Crampin and Pirani [14], de León and Rodrigues [27], Krupkova [25], Miron and Anastasiei [32]. For the differential geometry of a second order vector field one has to study its associated nonlinear connection and dynamical covariant derivative. Such nonlinear connection was introduced by Crampin [13] and Grifone [21] and it is a very important tool for the geometry of Finsler and Lagrange spaces developed by Miron and Anastasiei [32]. For a regular Lagrangian, the canonical nonlinear connection is the only connection that is metric and compatible with the symplectic structure, as it has been shown in [7].

Tangent bundles of higher order are canonically endowed with tangent structures of higher order, introduced by Eliopoulos [19]. This tangent geometry of higher order provides a special differential calculus, which has been investigated by many authors, for example by Yano and Ishihara [38], Tulczyjew [37], de León and Rodrigues [26], Saunders [34], Miron [31]. Important works that connect the geometry of higher-order differential equations with the geometry of higher order Lagrangian mechanics are due to Crampin et al. [15], de León and Rodrigues [28], Byrnes [9].

Spaces involving metric tensors whose components are functions of position and higher order accelerations were introduced first by Kawaguchi [22]. The variational problem of such metric structures of order k , $k \geq 1$, was investigated by Craig [12] and Synge [36] who associated to it a number of $k + 1$ covariant vectors. Among these vectors, only the last two live on the tangent space of order k . For a regular Lagrangian of order k , the last one determines one of the Cartan-Poincaré one forms, the one before the last one determines a vector field of order $k + 1$. This vector field was called by Miron in [30] the Craig-Synge vector field, or the canonical semispray, of the Lagrange space of order k . The geometry of higher order Lagrangians based on the associated semispray and the associated nonlinear connection has been developed recently by Miron [31]. From a semispray that lives on the tangent bundle of order k , one can derive different nonlinear connections as it has been done by Catz [11], de León and Rodrigues [26], Byrnes [9], Miron and Atanasiu [32], Bucataru [6]. For $k > 1$ these nonlinear connections are different and the expression of their coefficients is quite complicated. Geometric invariants induced by two different nonlinear connections associated to such a semispray were studied by Crampin and Saunders [16]. In this paper we limit our work to the case of second order Lagrangians.

A dynamical system on the total space of second order acceleration bundle T^2M is defined as third order vector field, S , which lives on T^2M and is mapped by the second order tangent map into one of the Liouville vector field. Such vector field is called a semispray. For a regular Lagrangian of second order, we prove that one can associate to it two Cartan-Poincaré one-forms and two Cartan-Poincaré two-forms. We show that one of the Cartan-Poincaré two-forms is a presymplectic form. Moreover, both vertical distributions are Lagrangian subbundles with respect to this presymplectic form.

The classical variational problem for a second order Lagrangian leads to fourth-order equations, which are the Euler-Lagrange equations. To the fourth-order Euler-Lagrange vector field, Craig [12] and Synge [36] associated another vector field of third order, which was called the Craig-Synge vector by Miron [30]. In this paper we study Craig-Synge equations and the corresponding vector field since they live on the second order tangent bundle of the base manifold. For this vector field, which is a semispray of the considered regular Lagrangian of second order, we prove that it is uniquely determined by the two associated Cartan-Poincaré one-forms. This result extends the work of Sarlet et al. [35] for the canonical second order vector field of a first order regular Lagrangian. We prove that the Craig-Synge covector can be obtained directly by a special variational principle of a second order regular Lagrangian for the case when only the vertical part of the curve is varied. It is known that for a Lagrangian of order greater than one, the integral of action along a curve does not depend on the parametrization of the curve only if the Lagrangian is degenerate. This is known as Zermelo condition, geometric aspects of this condition were investigated by Kondo, [23]. Therefore for a regular Lagrangian, when studying the integral of action, the parametrization of a curve is essential. The variation of the vertical part of the curve allows for variation of such parametrization. Recently, seismic ray path through higher order metric spaces, where the Zermelo condition is satisfied, were investigated by Yajima and Nagahama [39].

A nonlinear connection on the second order tangent bundle together with a semispray define dynamical covariant derivatives of first and second order. These covariant derivatives allow for invariant expressions of various geometric objects derived from a second order Lagrangian. In this paper we give conditions that uniquely determine a new nonlinear connection of a second order regular Lagrangian. One of these conditions states that the corresponding horizontal distribution is a Lagrangian subbundle for the presymplectic form. With respect to this nonlinear connection, the second order dynamical covariant derivative of the metric tensor vanishes. This generalizes the result for a first order regular Lagrangian, where the dynamical covariant derivative of the metric tensor vanishes, [7].

2. SECOND ORDER TANGENT BUNDLE

The space of second order accelerations can be identified with the space of second order tangent bundle. The space of higher order tangent bundles using jet theory was introduced by Ehresmann [18]. In this section we present some of the geometric structures that live on the total space of the second order tangent bundle, such as second order tangent structures, Liouville vector fields, vertical and horizontal distributions and semispray. These structures were investigated previously by Eliopoulos [19], Yano and Ishihara [38], Tulczyjew [37], de León and Rodrigues [26], Crampin et al. [15] and Miron [31].

For a real, n -dimensional, smooth manifold M , we denote by (T^2M, π^2, M) its tangent bundle of second order. A local chart $(U, \varphi = (x^i))$ on M induces a local chart $((\pi^2)^{-1}(U), \Phi = (x^i, y^{(1)i}, y^{(2)i}))$ on T^2M , where for a two-jet $j_0^2\rho \in T^2M$, the coordinate functions are defined as follows

$$y^{(\alpha)i}(j_0^2\rho) = \frac{1}{\alpha!} \frac{d^\alpha (x^i \circ \rho)}{dt^\alpha} \Big|_{t=0}, \quad \alpha \in \{1, 2\}.$$

The second order tangent structure J and second order cotangent structure J^* , which are called also vertical endomorphisms, are defined as follows

$$J = \frac{\partial}{\partial y^{(1)i}} \otimes dx^i + \frac{\partial}{\partial y^{(2)i}} \otimes dy^{(1)i}, \quad J^* = dx^i \otimes \frac{\partial}{\partial y^{(1)i}} + dy^{(1)i} \otimes \frac{\partial}{\partial y^{(2)i}}.$$

The foliated structure of T^2M allows for two regular vertical distributions, $V_1(u) = \text{Ker } d_u\pi^2 = \text{Ker } J^2 = \text{Im } J$ and $V_2(u) = J(V_1(u)) = \text{Ker } d_u\pi_1^2 = \text{Ker } J = \text{Im } J^2$, $\forall u \in T^2M$. These two distributions are integrable, the first one is tangent to the fibers of $\pi^2 : (x, y^{(1)}, y^{(2)}) \in T^2M \mapsto (x) \in M$, while the second one is tangent to the fibers of $\pi_1^2 : (x, y^{(1)}, y^{(2)}) \in T^2M \mapsto (x, y^{(1)}) \in TM$.

The following two vertical vector fields are globally defined on T^2M and they are called Liouville vector fields.

$$\mathbb{C}_2 = y^{(1)i} \frac{\partial}{\partial y^{(1)i}} + 2y^{(2)i} \frac{\partial}{\partial y^{(2)i}}, \quad \mathbb{C}_1 = J(\mathbb{C}_2) = y^{(1)i} \frac{\partial}{\partial y^{(2)i}}.$$

A semispray is a globally defined vector field S on T^2M that satisfies the equation $JS = \mathbb{C}_2$. Therefore, a semispray S , which is a third order vector field, can be expressed as follows

$$(1) \quad S = y^{(1)i} \frac{\partial}{\partial x^i} + 2y^{(2)i} \frac{\partial}{\partial y^{(1)i}} - 3G^i(x, y^{(1)}, y^{(2)}) \frac{\partial}{\partial y^{(2)i}}$$

and it is perfectly determined by its coefficient functions $G^i(x, y^{(1)}, y^{(2)})$. In this paper we provide conditions that uniquely determine such a semispray for a regular second order Lagrange space.

A nonlinear connection, or a horizontal distribution, on T^2M is a regular distribution $N : u \in T^2M \mapsto N(u) \subset T_u T^2M$ that is supplementary to the vertical distribution V_1 . In other words, the following direct sum holds true.

$$(2) \quad T_u T^2M = N(u) \oplus V_1(u), \forall u \in T^2M.$$

Since $d_u \pi^2 : T_u T^2M \mapsto T_{\pi^2(u)}M$ is an epimorphism of vector spaces, whose kernel is $V_1(u)$, its restriction to $N(u)$ is an isomorphism of vector spaces. We denote by $l_{h,u} : T_{\pi^2(u)}M \longrightarrow N(u)$ its inverse, to which we refer to as the horizontal lift induced by the nonlinear connection N . If we consider the regular, n -dimensional distribution $N_1(u) = J(N(u))$ then we have the following direct sums.

$$(3) \quad T_u T^2M = N(u) \oplus N_1(u) \oplus V_2(u), \quad V_1(u) = N_1(u) \oplus V_2(u), \quad \forall u \in T^2M.$$

An adapted basis to the first decomposition (3) is given by

$$(4) \quad \left. \frac{\delta}{\delta x^i} \right|_u := l_{h,u} \left(\left. \frac{\partial}{\partial x^i} \right|_{\pi^2(u)} \right), \quad \left. \frac{\delta}{\delta y^{(1)i}} \right|_u := J \left(\left. \frac{\delta}{\delta x^i} \right|_u \right), \quad \left. \frac{\partial}{\partial y^{(2)i}} \right|_u.$$

With respect to the natural basis of $T_u T^2M$, we use the following notations:

$$(5) \quad \left. \frac{\delta}{\delta x^i} \right|_u = \left. \frac{\partial}{\partial x^i} \right|_u - N_{(1)}^j(u) \left. \frac{\partial}{\partial y^{(1)j}} \right|_u - N_{(2)}^j(u) \left. \frac{\partial}{\partial y^{(2)j}} \right|_u.$$

Functions $N_{(1)}^j$ and $N_{(2)}^j$ are called local coefficients of the nonlinear connection N .

The dual basis of the adapted basis given by expression (4) is given by

$$(6) \quad dx^i, \quad \delta y^{(1)i} = dy^{(1)i} + M_{(1)}^i dx^j, \quad \delta y^{(2)i} = dy^{(2)i} + M_{(1)}^i dy^{(1)j} + M_{(2)}^i dx^j.$$

Functions $M_{(1)}^i$ and $M_{(2)}^i$ are called the dual coefficients of the nonlinear connection

and they are related to the coefficients of the nonlinear connection through the following formulas

$$(7) \quad M_{(1)}^i = N_{(1)}^i, \quad M_{(2)}^i = N_{(2)}^i + N_{(1)}^m N_{(1)}^m.$$

A semispray induces a nonlinear connection, which can be expressed in terms of some product structures as it has been shown by Catz [11] and de León and Rodrigues [26]. Different sets of dual coefficients of nonlinear connections induced by a semispray were obtained by Miron [31] and Bucataru [6]. In this paper we provide conditions that uniquely determine a nonlinear connection in terms of its compatibility with the metric structure and the presymplectic structure of a second order Lagrange space.

3. SECOND ORDER LAGRANGE SPACE

Important geometric structures for the geometry of a regular first order Lagrangian can be derived from the associated Cartan-Poincaré one and two-forms: Euler-Lagrange vector field, nonlinear connection, symmetries. In this section we introduce some of the geometric structures one can associate to a second order regular Lagrangian, such as Cartan-Poincaré one and two-forms. We study their

relations with some other geometric structures that live on the second order tangent bundle.

Consider $L : T^2M \longrightarrow \mathbb{R}$ a regular Lagrangian of second order. In other words, the metric tensor

$$(8) \quad g_{ij} = \frac{1}{2} \frac{\partial^2 L}{\partial y^{(2)i} \partial y^{(2)j}}$$

is a symmetric second order tensor field that has maximal rank n on T^2M .

A tensor field on T^2M is called a d -tensor field if its components change as the components of a similar tensor field on the base manifold, under corresponding change of coordinates. The metric tensor given by expression (8) is a second order d -tensor field since its components g_{ij} behave as the components of a $(0,2)$ -type tensor field on the base manifold.

For a regular Lagrangian of second order, one can define the following globally defined two Cartan-Poincaré one-forms:

$$(9) \quad \theta_L^1 = J^*(dL) = d_J L = \frac{\partial L}{\partial y^{(1)i}} dx^i + \frac{\partial L}{\partial y^{(2)i}} dy^{(1)i} \quad \text{and}$$

$$(10) \quad \theta_L^2 = (J^*)^2(dL) = d_{J^2} L = \frac{\partial L}{\partial y^{(2)i}} dx^i.$$

We consider also the following Cartan-Poincaré two-forms:

$$(11) \quad \begin{aligned} \omega_L^2 &= d\theta_L^2 = d\left(\frac{\partial L}{\partial y^{(2)i}} dx^i\right) = d\left(\frac{\partial L}{\partial y^{(2)i}}\right) \wedge dx^i = \\ &\frac{\partial^2 L}{\partial x^j \partial y^{(2)i}} dx^j \wedge dx^i + \frac{\partial^2 L}{\partial y^{(1)j} \partial y^{(2)i}} dy^{(1)j} \wedge dx^i + 2g_{ji} dy^{(2)j} \wedge dx^i, \end{aligned}$$

$$(12) \quad \omega_L^1 = d\theta_L^1 = d\left(\frac{\partial L}{\partial y^{(1)i}}\right) \wedge dx^i + d\left(\frac{\partial L}{\partial y^{(2)i}}\right) \wedge dy^{(1)i}.$$

We remark here that the regularity of the Lagrangian L implies the fact that $\text{rank}(\omega_L^2) = 2n < 3n = \dim(T^2M)$. We refer to ω_L^2 as to the canonical presymplectic structure of the Lagrangian L . These aspects were briefly discussed in [8].

Proposition 3.1. *Consider S an arbitrary semispray on T^2M . For a second order Lagrangian L , its Cartan-Poincaré one and two-forms have the following properties.*

- (1) $d_J \theta_L^2 = d_J d_{J^2} L = 0$, $d_{J^2} \theta_L^1 = d_{J^2} d_J L = 0$,
- (2) $i_S \theta_L^1 = \theta_L^1(S) = \mathbb{C}_2(L)$, $i_S \theta_L^2 = \theta_L^2(S) = \mathbb{C}_1(L)$,
- (3) $\mathcal{L}_S \theta_L^1 = i_S \omega_L^1 + d(\mathbb{C}_2(L))$, $\mathcal{L}_S \theta_L^2 = i_S \omega_L^2 + d(\mathbb{C}_1(L))$.

Proof. First two properties follow directly from expressions (11) and (12), if we compose with J and J^2 respectively. Next two properties are direct consequences of expressions (9) and (10). Last two properties follow from the previous ones if we take the exterior differential and use the fact that $\mathcal{L}_S = di_S + i_S d$, where $\mathcal{L}_S$ is the Lie derivative along the semispray S . $\square$

First property of Proposition 3.1 means that $\omega_L^2(JX, JY) = \omega_L^2(J^2X, J^2Y) = 0$, $\forall X, Y \in \chi(T^2M)$ and therefore both vertical distributions V_1 and V_2 are Lagrangian subbundles for the presymplectic structure ω_L^2 . Second property of Proposition 3.1 means that $\omega_L^1(J^2X, J^2Y) = 0$, $\forall X, Y \in \chi(T^2M)$ and therefore the vertical distribution V_2 is a Lagrangian subbundle for the Cartan-Poincaré two-form ω_L^1 .

We will explain now how the results in this section extend the well-known results for the first order case, to which we refer to Crampin and Pirani [14]. Consider $L : (x, y) \in TM \mapsto L(x, y) \in \mathbb{R}$ a first order regular Lagrangian and let

$$(13) \quad \theta_L = d_J L = \frac{\partial L}{\partial y^i} dx^i, \quad \omega_L = dd_J L = d \left(\frac{\partial L}{\partial y^i} \right) \wedge dx^i$$

be the Cartan-Poincaré one and two-forms. For these we have the following properties.

The vertical distribution of the tangent bundle is a Lagrangian subbundle for the symplectic structure ω_L . This is due to the fact that $d_J \theta_L = d_J^2 L = 0$ and therefore $\omega_L(JX, JY) = 0$, for all X, Y vector fields on TM .

For an arbitrary semispray S on TM , which is a second order vector field on the base manifold M , we have $i_S \theta_L = \mathbb{C}(L)$, where $\mathbb{C} = y^i (\partial / \partial y^i)$ is the Liouville vector field on TM . By taking the exterior differential of the above formula we obtain $i_S \omega_L = \mathcal{L}_S \theta_L - d(\mathbb{C}(L))$. For the Euler-Lagrange vector field S we have $\mathcal{L}_S \theta_L = dL$. Therefore, the equation $i_S \omega_L = -d(L - \mathbb{C}(L))$ uniquely determines the canonical semispray and implies that S is a Cartan symmetry for the Euler-Lagrange equations.

4. THIRD ORDER VECTOR FIELDS AND CRAIG-SYNGE COVECTORS

Spaces with metric structures whose components depend on higher order accelerations were introduced by Kawaguchi, [22]. The classic variational problem of a metric structure of order two induces the Euler-Lagrange covector field, which is of order four. From the corresponding vector field, which lives on the tangent bundle of order three, it is difficult to derive geometric objects that live on the tangent bundle of second order and to develop a geometric theory of the given Lagrangian of second order.

For a second order regular Lagrangian, using the Euler-Lagrange covector field, Craig [12] and Synge [36] derived another covector of order three along a curve. Miron [30] paid a special attention to this covector that was called Craig-Synge covector. The corresponding vector field lives on the tangent bundle of order two, it is a semispray and one can develop from it the geometry of the given Lagrangian.

In this section we prove, that the Craig-Synge covector can be independently derived using a variational principle of a regular second order Lagrangian, for the case when only the vertical components of the curve are varied. The corresponding vector field is a semispray associated to the second order Lagrange space. In this section we provide the equation that uniquely determines this canonical semispray of a second order regular Lagrangian. This equation generalizes the well-known equation for a first order Lagrangian.

Consider the following variational problem for the regular Lagrangian L . Let

$$c_\varepsilon : t \in [0, 1] \mapsto c_\varepsilon(t) = \left(x^i(t), \frac{dx^i}{dt}(t) + \varepsilon V^i(t), \frac{1}{2} \frac{d^2 x^i}{dt^2}(t) + \varepsilon \frac{dV^i}{dt}(t) \right) \in T^2M,$$

be a variation of the curve $c(t) = c_0(t)$ in T^2M , where ε belongs to some small neighborhood of $0 \in \mathbb{R}$, and $V^i(t)$ are the components of a vector field along curve c such that $V^i(0) = V^i(1) = 0$. We look for necessary conditions for the curve $c = c_0$ to be an extremal of the integral

$$I(c_\varepsilon) = \int_0^1 L\left(x^i(t), \frac{dx^i}{dt}(t) + \varepsilon V^i(t), \frac{1}{2} \frac{d^2 x^i}{dt^2}(t) + \varepsilon \frac{dV^i}{dt}(t)\right) dt.$$

For this we require that c is a solution of the following equation

$$(14) \quad \left. \frac{d}{d\varepsilon} (I(c_\varepsilon)) \right|_{\varepsilon=0} = 0.$$

Equation (14) can be written as follow

$$\begin{aligned} 0 &= \left. \frac{d}{d\varepsilon} (I(c_\varepsilon)) \right|_{\varepsilon=0} = \int_0^1 \left(\frac{\partial L}{\partial y^{(1)i}} V^i + \frac{\partial L}{\partial y^{(2)i}} \frac{dV^i}{dt} \right) dt \\ &= \int_0^1 \left[\frac{\partial L}{\partial y^{(1)i}} - \frac{d}{dt} \left(\frac{\partial L}{\partial y^{(2)i}} \right) \right] V^i dt + \left(\frac{\partial L}{\partial y^{(2)i}} V^i \right) \Big|_{t=0}^{t=1}. \end{aligned}$$

Since V is an arbitrary vector field, we have that equation (14) holds true if and only if the following Craig-Synge equations, [12], [36] are satisfied:

$$(15) \quad \frac{\partial L}{\partial y^{(1)i}} - \frac{d}{dt} \left(\frac{\partial L}{\partial y^{(2)i}} \right) = 0.$$

For a regular Lagrangian L , equations (15) represent a system of third order differential equations, which can be written as follows:

$$(16) \quad \frac{d^3 x^i}{dt^3} + 3! G^i \left(x, \frac{dx}{dt}, \frac{1}{2!} \frac{d^2 x}{dt^2} \right) = 0.$$

The functions G^i are given by

$$(17) \quad 3G^j = \frac{1}{2} g^{ji} \left[d_T \left(\frac{\partial L}{\partial y^{(2)i}} \right) - \frac{\partial L}{\partial y^{(1)i}} \right]$$

and they are local coefficients of a semispray, as it has been shown by Miron in [30]. We refer to the vector field S given by expression (1), whose coefficients are given by expression (17), as to the semispray S of the regular Lagrangian L . In expression (17), d_T is the Tulczyjew operator, [37]

$$d_T = y^{(1)i} \frac{\partial}{\partial x^i} + 2y^{(2)i} \frac{\partial}{\partial y^{(1)i}}.$$

Next theorem gives the equation that uniquely determines the semispray of a second order regular Lagrangian.

Theorem 4.1. *For a regular second order Lagrangian L , the semispray, whose coefficients are given by expression (17), is the only semispray S on T^2M that satisfies the following equation*

$$(18) \quad \mathcal{L}_S (d_{J^2} L) = d_J L.$$

Proof. Consider S a semispray on T^2M , given by expression (1). The Lie derivative of the Cartan-Poincaré one-form $\theta_L^2 = d_{J^2} L$ has the following expression

$$(19) \quad \mathcal{L}_S \theta_L^2 = \left[d_T \left(\frac{\partial L}{\partial y^{(2)i}} \right) - 6g_{ji} G^j \right] dx^i + \frac{\partial L}{\partial y^{(2)i}} dy^{(1)i}.$$

Using expression (19), we obtain that for a regular Lagrangian, equation (18) uniquely determines the functions G^i , which are given by expression (17). $\square$

Corollary 4.2. *For a regular second order Lagrangian L , its semispray is uniquely determined by the following equation*

$$(20) \quad i_S \omega_L^2 = -d(\mathbb{C}_1(L)) + \theta_L^1.$$

Proof. From Proposition 3.1 it follows that $\mathcal{L}_S \theta_L^2 = i_S \omega_L^2 + d(\mathbb{C}_1(L))$. Therefore, equations (18) and (20) are equivalent and each of them, uniquely determine the semispray of the regular, second order Lagrange space. $\square$

For a first order regular Lagrangian $L(x, y)$, as it has been shown by Sarlet et al. [35], the canonical semispray is uniquely determined by the following equation

$$(21) \quad \mathcal{L}_S(d_J L) = dL.$$

For a first order Lagrangian the corresponding equation for (20) reads as follows

$$(22) \quad i_S \omega_L = -d(\mathbb{C}(L) - L).$$

We remark here that in equations (21) and (22) the right hand-side is an exact form, while in equations (18) and (20) the right hand-side is not an exact form any more since θ_L^1 is not even a closed one-form. This has the following consequences, which are different from the case of a first order Lagrangian.

Corollary 4.3. *For a regular second order Lagrangian, its semispray satisfies:*

- (1) $\mathcal{L}_S \omega_L^2 = \omega_L^1$,
- (2) $S(\mathbb{C}_1(L)) = \mathbb{C}_2(L)$.

Proof. If we take the exterior differential of both sides of equation (18) and use the fact that the Lie derivative commutes with the exterior differential, we obtain $\mathcal{L}_S \omega_L^2 = \omega_L^1$. Using the skew symmetry of ω_L^2 and expression (20), we obtain $S(\mathbb{C}_1(L)) = \mathbb{C}_2(L)$. $\square$

From the above results we can see that the semispray S is not a symmetry for the corresponding second order Lagrange space.

5. SECOND ORDER DYNAMICAL DERIVATIVE

In this section, we introduce dynamical covariant derivatives of first and second order, induced by a pair (S, N) , where S is the associated semispray of a second order Lagrange space and N is a nonlinear connection. Using these dynamical derivatives and the presymplectic structure we give conditions that uniquely determine a nonlinear connection N . With respect to this nonlinear connection, the second order dynamical covariant derivative of the metric tensor vanishes and the horizontal distribution is a Lagrangian subbundle for the presymplectic structure.

Let S be the associated semispray of a second order Lagrange space, which is uniquely determined by Theorem 4.1 and consider N a nonlinear connection. For a d -vector field with components X^i , we define its first and second order dynamical derivatives as follows

$$(23) \quad \nabla X^i = S(X^i) + \underset{(1)}{M_j^i} X^j, \quad \nabla^2 X^i = S^2(X^i) + \underset{(1)}{2M_j^i} S(X^j) + \underset{(2)}{2M_j^i} X^j.$$

Remark. Let $X = X^i (\partial/\partial x^i)$ be a vector field on the base manifold M . Its complete lift, given by

$$(24) \quad X^c = X^i \frac{\partial}{\partial x^i} + S(X^i) \frac{\partial}{\partial y^{(1)i}} + \frac{1}{2} S^2(X^i) \frac{\partial}{\partial y^{(2)i}},$$

can be expressed as follows

$$(25) \quad X^c = X^i \frac{\delta}{\delta x^i} + \nabla X^i \frac{\delta}{\delta y^{(1)i}} + \frac{1}{2} \nabla^2 X^i \frac{\delta}{\delta y^{(2)i}}.$$

One can extend these first and second order dynamical derivatives to arbitrary d -tensor fields. In this paper we are interested in the first and second derivatives of the metric tensor g_{ij} . These derivatives are given by

$$(26) \quad \nabla g_{ij} = S(g_{ij}) - N_{(1)i}^k g_{kj} - N_{(1)j}^k g_{ik},$$

$$(27) \quad \begin{aligned} \nabla^2 g_{ij} &= S^2(g_{ij}) - 2N_{(1)i}^k S(g_{kj}) - 2N_{(1)j}^k S(g_{ik}) \\ &\quad - 2N_{(2)i}^k g_{kj} - 2N_{(2)j}^k g_{ik} + 2g_{mk} N_{(1)j}^k N_{(1)i}^m \end{aligned}$$

Consider $g = g_{ij} dx^i \otimes dx^j$ the metric tensor of a second order regular Lagrangian. One can extend this tensor to the following metric structure on T^2M

$$(28) \quad \begin{aligned} g^c &= g_{ij} \left(dx^i \otimes dy^{(2)j} + dy^{(1)i} \otimes dy^{(1)j} + dy^{(2)i} \otimes dx^j \right) + \\ &\quad S(g_{ij}) \left(dx^i \otimes dy^{(1)j} + dy^{(1)i} \otimes dx^j \right) + \frac{1}{2} S^2(g_{ij}) dx^i \otimes dx^j. \end{aligned}$$

With respect to the adapted basis of the nonlinear connection N , the metric structure g^c has the following expression

$$(29) \quad \begin{aligned} g^c &= g_{ij} \left(dx^i \otimes \delta y^{(2)j} + \delta y^{(1)i} \otimes \delta y^{(1)j} + \delta y^{(2)i} \otimes dx^j \right) + \\ &\quad \nabla g_{ij} \left(dx^i \otimes \delta y^{(1)j} + \delta y^{(1)i} \otimes dx^j \right) + \frac{1}{2} \nabla^2 g_{ij} dx^i \otimes dx^j. \end{aligned}$$

Theorem 5.1. *Let S be the associated semispray of a second order regular Lagrangian. There exists a unique nonlinear connection N on T^2M such that*

$$(1) \quad \omega_L^2(hX, v_1 Y) = 2g^c(hX, v_1 Y), \quad \forall X, Y \in \chi(T^2M).$$

$$(2) \quad \nabla^2 g = 0, \quad d_h \theta_L^2 = 0.$$

Proof. We prove that the first condition of the theorem uniquely determines the first coefficients $N_{(1)i}^k$, while the second condition uniquely determines the second

coefficients $N_{(2)i}^k$ of a nonlinear connection. We use the metric tensor g_{ij} to raise and lower indices, therefore we denote

$$N_{(1)ij} := g_{ik} N_{(1)j}^k, \quad N_{(2)ij} := g_{ik} N_{(2)j}^k.$$

The presymplectic structure ω_L^2 can be expressed, with respect to the adapted basis of a nonlinear connection N , as follows

$$(30) \quad \omega_L^2 = \frac{\delta}{\delta x^i} \left(\frac{\partial L}{\partial y^{(2)j}} \right) dx^j \wedge dx^i + \frac{\delta}{\delta y^{(1)i}} \left(\frac{\partial L}{\partial y^{(2)j}} \right) \delta y^{(1)j} \wedge dx^i + 2g_{ij} \delta y^{(2)j} \wedge dx^i.$$

Condition 1 of the theorem can be written as follows

$$(31) \quad \omega_L^2 \left(\frac{\delta}{\delta x^i}, \frac{\delta}{\delta y^{(1)j}} \right) = \frac{\delta}{\delta y^{(1)i}} \left(\frac{\partial L}{\partial y^{(2)j}} \right) = 2g^c \left(\frac{\delta}{\delta x^i}, \frac{\delta}{\delta y^{(1)j}} \right) = 2g_{ij|},$$

which is equivalent to

$$(32) \quad 2N_{ij}^{(1)} + 4N_{ji}^{(1)} = 2S(g_{ij}) + \frac{\partial^2 L}{\partial y^{(1)i} \partial y^{(2)j}}.$$

Since $\nabla g_{ij} = g_{ij|}$ are the components of a second rank symmetric d -tensor field, from expression (31) we obtain the symmetry in i and j of the following expression

$$\frac{\delta}{\delta y^{(1)i}} \left(\frac{\partial L}{\partial y^{(2)j}} \right),$$

which is equivalent to

$$(33) \quad 2N_{ij}^{(1)} - 2N_{ji}^{(1)} = \frac{\partial^2 L}{\partial y^{(2)i} \partial y^{(1)j}} - \frac{\partial^2 L}{\partial y^{(2)j} \partial y^{(1)i}}.$$

From expressions (32) and (33) we obtain that $N_{ij}^{(1)}$ is uniquely determined and it has the following expression

$$(34) \quad N_{ij}^{(1)} = \frac{1}{3}S(g_{ij}) + \frac{1}{3} \frac{\partial^2 L}{\partial y^{(2)i} \partial y^{(1)j}} - \frac{1}{6} \frac{\partial^2 L}{\partial y^{(2)j} \partial y^{(1)i}}.$$

Using expression (27), we obtain that first part of condition 2 of the theorem, $\nabla^2 g_{ij} = 0$, uniquely determines the symmetric part of $N_{ij}^{(2)}$.

$$(35) \quad 2N_{ij}^{(2)} + 2N_{ji}^{(2)} = S^2(g_{ij}) - 2S(g_{ik})N_j^k - 2S(g_{kj})N_i^k + 2g_{mk}N_j^k N_i^m.$$

Second part of condition 2 of the theorem can be written as follows $\omega_L^2(hX, hY) = 0, \forall X, Y \in \chi(T^2M)$. From expression (30) this condition implies the symmetry in i and j of the following expression

$$\frac{\delta}{\delta x^i} \left(\frac{\partial L}{\partial y^{(2)j}} \right),$$

which determines the skewsymmetric part of $N_{ij}^{(2)}$.

$$(36) \quad 2N_{ij}^{(2)} - 2N_{ji}^{(2)} = \frac{\partial^2 L}{\partial y^{(2)i} \partial x^j} - \frac{\partial^2 L}{\partial y^{(2)j} \partial x^i} - N_j^k \frac{\partial^2 L}{\partial y^{(2)i} \partial y^{(1)k}} + N_i^k \frac{\partial^2 L}{\partial y^{(2)j} \partial y^{(1)k}}.$$

Using expressions (35) and (36), we obtain that the coefficients $N_{ij}^{(2)}$ are uniquely determined. $\square$

Corollary 5.2. *With respect to the nonlinear connection determined by Theorem 5.1, the presymplectic structure $\omega_L^2 = d\theta_L^2$ has the following form*

$$(37) \quad \omega_L^2 = d\theta_L^2 = 2g_{ij|}\delta y^{(1)j} \wedge dx^i + 2g_{ij}\delta y^{(2)j} \wedge dx^i.$$

If h , v_1 and v_2 are the projectors that correspond to the nonlinear connection determined by Theorem 5.1, then the Poincaré-Cartan one-form θ_L^2 satisfies the following equations

$$(38) \quad d_h \theta_L^2 = d_{v_1} \theta_L^2 = d_{v_2} \theta_L^2 = 0.$$

Proof. First condition and second part of the second condition of Theorem 5.1 are equivalent with the fact that the presymplectic structure ω_L^2 has with respect to adapted basis of the nonlinear connection the form given by expression (37). All equations (38) can be obtained directly from expression (37) if we compose it with the projectors h , v_1 and v_2 respectively. $\square$

Theorem 5.1 gives conditions that uniquely determine a nonlinear connection for a second order Lagrange space. These conditions can be restated as follows, there is a unique nonlinear connection such that $\nabla^2 g = 0$ and the presymplectic structure ω_L^2 is given by expression (37). However, the expressions we obtained for the coefficients $N_{ij}^{(1)}$ and $N_{ij}^{(2)}$ in the proof of the above theorem are not easy to use. Next proposition gives a simpler form for the first coefficients of the considered nonlinear connection.

Proposition 5.3. *Consider S the associated semispray of a second order regular Lagrangian L , whose coefficients G^i are given by expression (17). The following functions are the first coefficients of the nonlinear connection determined by Theorem 5.1.*

$$(39) \quad N_{(1)}^i = \frac{\partial G^i}{\partial y^{(2)j}},$$

Proof. We have to show that the functions $N_{(1)}^i$ given by expression (39) satisfy expression (34), which uniquely determine the first coefficients of the nonlinear connection. The associated semispray S of a second order regular Lagrangian has the local coefficients given by expression (17) and it is uniquely determined by the following condition

$$(40) \quad S \left(\frac{\partial L}{\partial y^{(2)i}} \right) = \frac{\partial L}{\partial y^{(1)i}}.$$

By direct calculation we have the following formula

$$(41) \quad \left[S, \frac{\partial}{\partial y^{(2)i}} \right] = -2 \frac{\partial}{\partial y^{(1)i}} + 3 \frac{\partial G^j}{\partial y^{(2)i}} \frac{\partial}{\partial y^{(2)j}}.$$

If we apply both sides of expression (41) to $\partial L / \partial y^{(2)k}$ and use expression (40) we obtain

$$(42) \quad g_{jk} \frac{\partial G^j}{\partial y^{(2)i}} = \frac{1}{3} S(g_{ik}) + \frac{1}{3} \frac{\partial^2 L}{\partial y^{(2)k} \partial y^{(1)i}} - \frac{1}{6} \frac{\partial^2 L}{\partial y^{(2)i} \partial y^{(1)k}}.$$

From expressions (34) and (42) we obtain that first coefficients $N_{(1)}^i$ of the nonlinear connection are given by expression (39). $\square$

According to Proposition 5.3 the coefficients $N_{(1)}^i$ of the nonlinear connection determined by Theorem 5.1 are the same for the nonlinear connections that were considered by Catz [11], Dodson and Radivoioci [17], de Leon and Rodrigues [26],

Byrnes [9], Miron [31] and Bucataru [6]. However, the other coefficients N_j^i of the nonlinear connection we obtained in Theorem 5.1 are different and with respect to them the presymplectic form has a very simple form given by expression (37).

6. EXAMPLES

In this section we consider examples of second order regular Lagrangians, which were previously used by Miron and Atanasiu [33] to extend semi-Riemannian, Finslerian and Lagrangian structures from the base manifold to the total space of second order tangent bundles. For such second order regular Lagrangians we compute the associated semispray and the nonlinear connection as they have been defined in the previous sections. If the Lagrangian function depends on position effectively, this nonlinear connection is different from the prolongation of a nonlinear connection introduced by Catz [11] and used by Miron and Atanasiu [33]. It is also different from the nonlinear connection used by Dodson and Radivoiović [17] and de León and Vázquez [26] to study the geometry of the tangent bundle of order two. Each of these nonlinear connections give information regarding the Lagrangian function as we will see in this section. However, for the semi-Riemannian case, with respect to the nonlinear connection introduced in the previous section, the first and second order covariant derivatives of the metric tensor vanish and the presymplectic structure has a very simple form with respect to it.

Let $g = g_{ij}(x) dx^i \otimes dx^j$ be a semi-Riemannian metric on the base manifold M and denote by $\gamma_{jk}^i(x)$ the Christoffel symbols of the Riemannian metric. Consider the first order Lagrangian $L_1 : TM \rightarrow \mathbb{R}$ given by

$$(43) \quad L_1(x, y^{(1)i}) = g_{ij}(x) y^{(1)i} y^{(1)j} = \|y^{(1)}\|^2.$$

We recall that the geodesics of the semi-Riemannian metric g , which are extremal curves for the first order Lagrangian L_1 , are solutions of the following system of equations:

$$(44) \quad \nabla \left(\frac{dx^i}{dt} \right) = 0.$$

Using the fact that $\nabla g = 0$, for a curve $c(t) = (x^i(t), y^{(1)i}(t))$ on TM we have that

$$(45) \quad \frac{d}{dt} \left(L_1(x(t), y^{(1)i}(t)) \right) = 2g_{ij}(x(t)) \frac{dx^i}{dt} \nabla y^{(1)i},$$

from which we obtain that L_1 is conserved along the geodesics of the semi-Riemannian metric g . The right hand side of expression (45) is the first-order deformation Lagrangian $\mathcal{L}_1$, considered by Casciaro and Fracavaglia in [10], expression (3.3).

By direct calculation it follows that

$$(46) \quad z^{(2)i} = y^{(2)i} + \frac{1}{2} \gamma_{jk}^i(x) y^{(1)j} y^{(1)k} = \frac{1}{2} \nabla y^{(1)i}$$

are the components of a d -vector field on T^2M , which represents the covariant expression of the second order acceleration. The function $L_2 : T^2M \rightarrow \mathbb{R}$, given by

$$(47) \quad L_2(x, y^{(1)}, y^{(2)}) = g_{ij}(x) z^{(2)i} z^{(2)j} = \frac{1}{4} \|\nabla y^{(1)i}\|^2$$

is well defined on the second order tangent space, it is a second order regular Lagrangian and represents the square of the magnitude of the second order acceleration. The following formulas can be obtained by a straightforward calculation.

$$(48) \quad \frac{\partial L_2}{\partial y^{(2)i}} = 2g_{ij}z^{(2)j}, \frac{\partial L_2}{\partial y^{(1)i}} = 2g_{kj}z^{(2)j}\gamma_{ip}^k y^{(1)p}, \frac{\partial^2 L_2}{\partial y^{(1)i}\partial y^{(2)j}} = 2g_{kj}\gamma_{ip}^k y^{(1)p}.$$

The associated semispray of the second order regular Lagrangian given by expression (47) is uniquely determined by Theorem 4.1. Its coefficients are given by

$$(49) \quad \begin{aligned} 3G^i &= d_T \left(z^{(2)i} \right) + \gamma_{jk}^i z^{(2)j} y^{(1)k} \\ &= \frac{1}{2} \left(\frac{\partial \gamma_{jk}^i}{\partial x^m} + \gamma_{pj}^i \gamma_{km}^p \right) y^{(1)j} y^{(1)k} y^{(1)m} + 3\gamma_{jk}^i y^{(1)j} y^{(2)k}. \end{aligned}$$

The corresponding nonlinear connection of the second order Lagrangian L_2 is uniquely determined by Theorem 5.1. First set of coefficients of this nonlinear connection are given by expression (39), which in our case are given by

$$(50) \quad N_{(1)}^i \left(x, y^{(1)} \right) = \frac{\partial G^i}{\partial y^{(2)j}} = \gamma_{jk}^i(x) y^{(1)k}$$

With respect to the nonlinear connection studied in [6] and using the second order covariant derivative, the Craig-Synge equations, given by expression (16), of the second order Lagrangian L_2 can be expressed in a very simple form, which generalizes the geodesic equation (44)

$$(51) \quad \nabla^2 \left(\frac{dx^i}{dt} \right) = 0.$$

With respect to the nonlinear connection introduced by Miron and Atanasiu in [33], for a curve $c(t) = (x^i(t), dx^i/dt)$ on TM we have the following formula

$$(52) \quad \frac{d^2}{dt^2} \left(L_1 \left(x, \frac{dx}{dt} \right) \right) = 8L_2 \left(x, \frac{dx}{dt}, \frac{1}{2} \frac{d^2 x}{dt^2} \right) + 2g_{ij}(x(t)) \frac{dx^i}{dt} \nabla^2 \left(\frac{dx^j}{dt} \right),$$

which is equivalent to

$$(53) \quad \frac{d^2}{dt^2} \left\| \frac{dx}{dt} \right\|^2 = 2 \left\| \nabla \left(\frac{dx^i}{dt} \right) \right\|^2 + 2g_{ij}(x(t)) \frac{dx^i}{dt} \nabla^2 \left(\frac{dx^j}{dt} \right).$$

With respect to the nonlinear connection introduced in the previous section the presymplectic structure and the complete lift have a very simple form. From expression (37) we obtain that the canonical presymplectic structure of the second order Lagrange space can be expressed as follows

$$(54) \quad \omega_{L_2}^2 = 2g_{ij} \delta y^{(2)j} \wedge dx^i.$$

From expression (29), the complete lift g^c of the Riemannian metric tensor can be expressed as follows:

$$(55) \quad g^c = g_{ij} \left(dx^i \otimes \delta y^{(2)j} + \delta y^{(1)i} \otimes \delta y^{(1)j} + \delta y^{(2)i} \otimes dx^j \right).$$

In this section we have seen that the covariant derivatives induced by the considered semispray and the nonlinear connection give simple covariant expressions for the geometric structures associated to the second order Lagrangian L_2 .

Acknowledgement. This work was supported by grants PN II IDEI 398 (I.B.) and by CEEEX 252 (R.M.) of the Romanian Ministry of Education.

REFERENCES

- [1] Abraham, R., Marsden, J.E.: *Foundation of Mechanics*. Benjamin/Cummings Publishing Company, 1978.
- [2] Ambrose, W., Palais, R.S., Singer, I.M.: *Sprays*. An. Acad. Brasil. **32**, 163–178 (1960).
- [3] Antonelli, P.L., Bucataru, I.: *Volterra-Hamilton production models with discounting: general theory and worked examples*. Nonlinear Analysis: Real World Applications. **2**, 337–356 (2001).
- [4] Arnold, V.I.: *Mathematical methods of classical mechanics*. Springer, 1978.
- [5] Anastasiei, M.: *Symplectic structures and Lagrange geometry*. In Anastasiei, M., Antonelli P.L. (eds) *Finsler and Lagrange Geometry*. Kluwer Academic Publisher, 9–16 (2003).
- [6] Bucataru, I.: *Sprays and homogeneous connections in the higher order geometry*. Stud. Cerc. Mat. **50** (5-6), 307–315 (1998).
- [7] Bucataru, I.: *Metric nonlinear connections*. Differential Geometry and its Application. **35** (3), 335–343 (2007).
- [8] Bucataru, I.: *Canonical semispray for higher order Lagrange spaces*. Comptes Rendus Mathématique Acad. Sci. Paris, Ser. I, **345**, 269–272 (2007).
- [9] Byrnes, G. B.: *A linear connection for higher-order ordinary differential equations*. Journal of Physics A. **29** (8), 1685–1694 (1996).
- [10] Casciaro, B., Francaviglia, M.: *A new variational characterization of Jacobi fields along geodesics*. Annali di Matematica pura ed applicata. **172** (1), 219–228 (1997).
- [11] Catz, G.: *Sur le fibré tangent d'ordre 2*. Comptes Rendus Acad. Sci. Paris. **278**, 178–182 (1974).
- [12] Craig, H.V.: *On a generalized tangent vector*. American Journal of Math. **57**, 457–462 (1935).
- [13] Crampin, M.: *On horizontal distributions on the tangent bundle of a differentiable manifold*. J. London Math. Soc. **2** (3), 178–182 (1971).
- [14] Crampin, M., Pirani, F.A.E.: *Applicable Differential Geometry*. Cambridge University Press. London Mathematical Society Lecture Notes, 59, 1986.
- [15] Crampin, M., Sarlet, W., Cantrijn, F.: *Higher order differential equations and higher order Lagrangian Mechanics*. Proc. Camb. Phil. Soc. **99**, 565–587 (1986).
- [16] Crampin, M., Saunders, D. J.: *On the geometry of higher-order ordinary differential equations and the Wuenschmann invariant*. Groups, geometry and physics, Monogr. Real Acad. Ci. Exact. Fis.-Quím. Nat. Zaragoza, 29, Acad. Cienc. Exact. Fis. Quím. Nat. Zaragoza, Zaragoza, 79–92 (2006).
- [17] Dodson, C.T.J., Radivoioci, M.: *Tangent and frame bundles of order two*. An. St. Univ. "Al.I. Cuza" Iasi. **28** (1), 63–71 (1982).
- [18] Ehresmann, C.: *Les prolongements d'une variété différentiable. I. Calcul des jets, prolongement principal*. Comptes Rendus Acad. Sci. Paris. **233**, 598–600 (1951).
- [19] Eliopoulos, H.A.: *On the general theory of differentiable manifolds with almost tangent structure*. Canad. Math. Bull. **8**, 721–748 (1965).
- [20] Godbillon, C.: *Geometrie Differentielle et Mecanique Analytique*. Hermann, Paris, 1969.
- [21] Grifone, J.: *Structure presque-tangente et connexions I*. Ann. Inst. Henri Poincaré. **22** (1), 287–334 (1972).
- [22] Kawaguchi, A.: *Die Differentialgeometrie in der verallgemeinerten Mannigfaltigkeit*. Rendiconti Circolo Matematico di Palermo. **56**, 245–276 (1932).
- [23] Kondo, K.: *On the physical meaning of the Kawaguchi space*. Tensor N.S. **14**, 191–215 (1963).
- [24] Klein, J.: *Espaces variationnels et mécanique*. Inst. Fourier (Grenoble). **12**, 1–124 (1962).
- [25] Krupkova, O.: *The geometry of ordinary variational equations*. Springer, 1997.
- [26] de León, M., Rodrigues, P. R.: *Generalized Classical Mechanics and Field Theory*. North-Holland Publishing Co., Amsterdam, 1985.
- [27] de León, M., Rodrigues, P. R.: *Methods of differential geometry in analytical mechanics*. North-Holland Mathematics Studies, 158. North-Holland Publishing Co., Amsterdam, 1989.
- [28] de León, M., Rodrigues, P. R.: *The inverse problem of Lagrangian dynamics for higher-order differential equations: a geometrical approach*. Inverse Problems. **8**(4), 525–540, (1992).
- [29] de León, M., Vasquez, E.: *On the geometry of the tangent bundle of order two*. An. Univ. Bucuresti. **34**, 40–48 (1985).

- [30] Miron, R.: *Noether theorem in higher order Lagrangian mechanics*. International Journal of theoretical Physics. **34** (7), 1123–1146.
- [31] Miron, R.: *The Geometry of Higher-Order Lagrange Spaces. Applications to Mechanics and Physics*. Kluwer Academic Publisher, FTPH no.82, 1997.
- [32] Miron, R., Anastasiei, M.: *The Geometry of Lagrange Spaces: Theory and Applications*. Kluwer Academic Publisher, FTPH no.59, 1994.
- [33] Miron, R., Atanasiu, Gh.: *Prolongation of Riemannian, Finslerian and Lagrangian structures*. Rev. Roumaine Math. Pures Appl. **41** (3-4), 237–249(1996).
- [34] Saunders, D.J.: *The geometry of jet bundles*. Cambridge university Press (1989).
- [35] Sarlet W., Cantrijn F., Crampin, M.: *A new look at second order equations and Lagrangian mechanics*. J. Phys. A: Math. Gen. **17**(10), 1999–2009 (1984).
- [36] Synge, J.L.: *Some intrinsic and derived vectors in a Kawaguchi space*. American Journal of Math. **57**, 679–691 (1935).
- [37] Tulczyjew, W.M.: *The Lagrange differential geometry*. Bull. Acad. Polon. Sci. **24**, 1089–1096 (1976).
- [38] Yano, K., Ishihara, S.: *Tangent and cotangent bundles*. Marcel Dekker Inc, 1976.
- [39] Yajima, T., Nagahama, H.: *Kawaguchi space, Zermelo's condition and seismic ray path*. Nonlinear Analysis: Real World Applications. **8** 130–135 (2007).

FACULTY OF MATHEMATICS, "AL.I.CUZA" UNIVERSITY, IASI, 700506, ROMANIA

E-mail address: bucataru@uaic.ro, radu.miron@uaic.ro

URL: <http://www.math.uaic.ro/~bucataru>